

Gas Phase Chromatography of the Bromides of Mo, Zr, and Nb

E. Sylwester, K. E. Gregorich, J. L. Adams, Y. H. Chung, M. R. Lane, C. A. Laue, D. M. Lee, C. A. McGrath, D. A. Shaughnessy, D. A. Strellis, P. A. Wilk, D. C. Hoffman

The Heavy Element Volatility Instrument (HEVI)¹ has been used to investigate the volatilities of Mo, Zr, and Nb bromides. HEVI is an on-line isothermal gas phase chromatography system which separates short lived isotopes based on the volatility of their molecular form. The 8-min ⁸⁸Mo, along with multiple isotopes of Nb and Zr, was produced at the LBNL 88-Inch Cyclotron via the reaction of 86 MeV ²⁸Si projectiles with a ^{nat}Cu target. Reaction products were transported by a He/KBr aerosol gas jet system and continuously collected on a quartz wool plug kept at 900° C in a quartz chromatography column. HBr was added at a rate of 150 ml/min in order to form the volatile metal-bromide species, which were then swept into the isothermal section of the chromatography column. The separated species were then reattached to KBr aerosols and transported through a capillary into a glass wool filter placed in front of an intrinsic Ge gamma detector. ⁸⁸Mo ($t_{1/2}$ 8.0 m), ⁸⁷Nb ($t_{1/2}$ 2.6 m), and ⁸⁵Zr ($t_{1/2}$ 7.9 m), were identified by gamma spectroscopy, and the volatilities of their brominated forms were investigated by measuring the yields of these isotopes simultaneously as a function of the temperature of the column. Figure 1 shows the volatility data collected for the most prominent gamma line of each isotope. Volatility temperature is defined here as the point at which 50% chemical yield through the column is observed for that species; at this point the length of time spent in the column is equal to the half-life of the isotope studied. The apparent yield at low temperatures for these species seen in Figure 1 has been studied previously and may be due to mechanical transport through the column by undecomposed aerosols³.

⁸⁷NbBr₅ was observed to be volatile at 150° C, and ⁸⁵ZrBr₄ was observed to be volatile at 250° C.

Preliminary data analyses shows no clear volatility temperature for the Mo species.

A Monte Carlo simulation program² was used to calculate the adsorption enthalpy of these species based on their volatilities. The adsorption enthalpy for NbBr₅ was calculated to be -100±5 kJ/mol. The adsorption enthalpy for ZrBr₄ was calculated to be -110±5 kJ/mol. Data analysis is still ongoing for Mo and so no adsorption enthalpy value has been calculated.

Footnotes and References

1. B. Kadkhadayan et al., *Nucl Instr. Meth.* **A317**, 254 (1992)
2. A. Türler, K. E. Gregorich, D. C. Hoffman, D. M. Lee, H. W. Gäggeler, Nuclear Science Division Annual Report, LBL-31855, p. 56 (1991).
3. E. Sylwester et al., Nuclear Science Division Annual Report, LBNL-39764, p. 61 (1995-96).




Fig. 1. Volatilities and calculated adsorption enthalpies for ⁸⁸MoBr₆, ⁸⁷NbBr₅, and ⁸⁵ZrBr₄. The Monte Carlo curves shown are the best fit to the data. No adsorption enthalpy curve has yet been calculated for Mo.